Full Paper

Registration Code of Publication: 11-28-19-16 Publication is available for discussion in the framework of the on-line Internet conference "Butlerov readings". http://butlerov.com/readings/ Contributed: October 27, 2011.

## **Development of methodology for the quantitative** Determination the amount of flavonoids in the flowers of Calendula officinalis of different sorts

© Konstantin I. Korbakov,<sup>1\*+</sup> Galina A. Tselikova,<sup>1</sup> Elena L. Malankina<sup>2</sup> and Lilia V. Kuznetsova<sup>2</sup>

<sup>1</sup>Department of Organic Chemistry and Chemical Dyes. A.N. Kosygin Moscow State Textile University. Malaya Kaluzhskaya St., 1. Moscow, GSP-1, 119071. Russia. *Phone:* +7 (495) 955-35-48. *E-mail: kobrakovk@mail.ru* <sup>2</sup> Department of Viticulture and Winemaking RSAU - K.A. Timiryasev MA. Timiryasev St., 49. block No. 17. Moscow, 127550. Russia. Phone: +7 (499) 976-49-06. E-mail: gandurina@mail.ru

\*Supervising author; <sup>+</sup>Corresponding author

Keywords: defining the sum of flavonoids, Calendula officinalis, natural dyes.

## Abstract

This paper describes the methodology for determining the amount of flavonoids in *Calendula officinalis* flowers by routine, which is based on the bathochromic shift of maximum absorption in the UV spectrum of the extract at the complexation of flavonoids with aluminum salts.