Full Paper Registration Code of Publication: 12-30-6-89 Publication is available for discussion in the framework of the on-line Internet conference "Butlerov readings". http://butlerov.com/readings/ Contributed: July 24, 2012.

Synthesis and properties of titanium-containing catalyst of hindered phenols oxidation

© Renat M. Akhmadullin,¹⁺ Dinar R. Gatiyatullin,¹ Alexander A. Momzyakov,¹ Svetlana I. Aghajanian,^{1*} Alfia G. Akhmadullina,² and Natalia A. Mukmeneva^{1*}

¹ Department of Synthetic Rubber Technology. Kazan National Research Technological University. K. Marx St., 68. Kazan, 420015. Tatarstan Republic. Russia. Phone: +7 (843) 231-42-13. *E-mail: ahmadullinr@gmail.com*

² Individual Entrepreneur. Yu. Fuchik St., 44-132. Kazan, 420139. Tatarstan Republic. Russia. Phone: +7 (843) 269-25-28. E-mail: ahmadullins@gmail.com

*Supervising author; ⁺Corresponding author

Keywords: oxidation, sterically hindered phenols, diphenoquinone, titanium oxide, alkaline catalysis.

Abstract

A process for producing a heterogeneous alkali catalyst for the oxidation of sterically hindered phenols has been proposed. The surface composition and stability of the proposed heterogeneous alkaline catalyst were studied. The principle of spatial hydrofobisation of alkaline catalyst surface was shown.