*Registration Code of Publication:* 12-32-12-112 Subsection: Physiogenic Biominerals. Publication is available for discussion in the framework of the on-line Internet conference "Butlerov readings". http://butlerov.com/readings/ Contributed: November 11, 2012.

## Thermodynamics calculations algorithm of the formation feasibility of sparingly soluble compounds of abiotic and biotic origin

© Olga A. Golovanova,<sup>1</sup>\*<sup>+</sup> Rinat R. Izmailov,<sup>1</sup> and Lyudmila V. Klimova<sup>2</sup>

<sup>1</sup>Department of Inorganic Chemistry. F.M. Dostoevsky Omsk State University. Pr. Mira St., 55a. Omsk, 644077. Omsk Region. Russia. Phone: +7 (3812) 64-27-00. E-mail: golovanoa2000@mail.ru <sup>2</sup>Department «Technology of Ceramics, Glass and Binders». South-Russian State Technical University. Prosvescheniya St., 132. Novocherkassk, 346428. Rostov Region. Russia. Phone: +7 (3812) 64-27-00. E-mail: izmailov 87@mail.ru

\*Supervising author; <sup>+</sup>Corresponding author

*Keywords:* thermodynamics, supersaturation index, Gibbs energy, formation of soluble compounds, biological fluids.

## Abstract

In the work, we offer the algorithm of thermodynamic calculations allowing on the basis of computed criteria to predict the feasibility of the formation of different mineralization in the human body. It is shown that this approach is consistent with experimental data on the phase composition of renal, dental and salivary stones in the human body.