Reference Object Identifier – ROI: jbc-01/18-53-1-143 The Digital Object Identifier - DOI: 10.37952/ROI-jbc-01/18-53-1-143 Submitted on January 05, 2018.

## The interaction of methyl pheophorbide *a* with diethanolamine

© Lyudmila A. Tulaeva,<sup>1+</sup> Elena A. Kustysheva,<sup>1</sup> and Dmitry V. Belykh<sup>2\*</sup>

<sup>1</sup>Department of Chemistry. Institute of Natural Sciences. Syktyvkar State University. Orktyabrsky Ave., 55. Syktyvkar, 167000. Russia. E-mail: tulaeva65@mail.ru <sup>2</sup>Laboratory of Organic Synthesis and Chemistry of Natural Compounds. Institute of Chemistry, Komi Scientific Center, Ural Branch. Russian Academy of Sciences. Pervomayskava St., 48. Syktyvkar, 167000. Russia. E-mail: belykh-dv@mail.ru

\*Supervising author; <sup>+</sup>Corresponding author

*Keywords:* methylpheophorbide *a*, chlorine e<sub>6</sub> amides, exocycle of methylpheophorbide *a*, diethanolamine, monoethanolamine, purpurin 18, purpurin 18 N-hydroxyethylimide, nucleophilic substitution reaction

## Abstract

At present work, the interaction of methyl pheophorbide a with diethanolamine was studied in order to synthesize the 13-amide chlorin e<sub>6</sub> derivative with two hydroxyl groups in the molecule to carry out the nucleophilic opening reaction of the exocycle E. This reaction does not require additional activation of the starting compound, and it usually occurs in mild conditions and with good yields. Carrying out such a reaction with diethanolamine containing two hydroxyl groups allowed the introduction of two hydroxyl groups into the molecule of the 13-amide chlorin e6 derivative in one step. The presence of two hydroxyl groups in the compound can promote the overall hydrophilicity of the corresponding derivative, improve the solubility in water and physiological fluids when used. The interaction of methyl pheophorbide a with diethanolamine was studied in chloroform and tetrahydrofuran at room temperature. The formation of new compounds occurs only when the reaction is carried out in chloroform, in the case of tetrahydrofuran, the formation of new compounds has not been observed. When the reaction was carried out in chloroform, the yield of the desired compound – chlorin e<sub>6</sub> 13-*N*,*N*-di-(2-hydroxyethylamide)-15,17-dimethyl ether – was 10%. It was found that, in addition to the target compound, a side product, N-(2-hydroxyethyl) inide of the purpurin methyl ester 18, was formed. The formation of N-hydroxyethylimide purpurin 18 appears to occur when monoethanolamine, present in small amounts in diethanolamine, is added to the purpurin methyl ester 18, which in turn can be formed by oxidation of methyl pheophorbide with air oxygen in the presence of a base. Disclosure of the exocycle E in methyl pheophorbide a and the formation of the main product of the reaction tertiary amine 13-N,N-di-(2-hydroxyethylamide)-15,17-dimethyl chlorin e6 and the side product of the reaction of Nhydroxyethylimide purpurin 18 was confirmed by IR, electronic (UV-Vis), and NMR spectroscopy. According to NMR 13-N,N-di-(2-hydroxyethylamide)-15,17-dimethyl ester of chlorin e<sub>6</sub>, like all tertiary 13amides of chlorin e<sub>6</sub>, forms two atropisomers differing from each other by the mutual arrangement of the planes of the amide group and a chlorine macrocycle.

## References

- [1] Chemical transformations of bacteriochlorophyll a and its medical applications M.A. Grin and A.F. Mironov. Russian Chemical Bulletin, International Edition. 2016. Vol.65. No.2. P.333-349.
- [2] O.I. Gushchina, M.Yu. Grishina, E.A. Larkina, V.S. Lebedeva, and A.F. Mironov. Synthesis of Hydroxy. Derivatives of Chlorin e6 Macroheterocycles. 2017. Vol.10. No.1. P.81-83. (russian)
- [3] D.V. Belykh, I.S. Tarabukina, I.V. Gruzdev, A.V. Kuchin. Aminomethylation of chlorophyll a phorbine derivatives using bis(N,N-dimethylamino)methane. Macroheterocycles. 2010. Vol.3. No.2-3. P.145-149.
- [4] D.V. Belykh. Isolation of chlorophyll derivatives from plant raw materials and synthesis of chlorine amides e6. PhD thesis in the chem. sciences. Syktyvkar. 2001. (russian)
- [5] D.V. Belykh, L.P. Karmanova, L.V. Spirikhin, and A.V. Kuchin. Synthesis of Amide Derivatives of Chlorin e6. Russian Journal of Organic Chemistry. 2007. Vol.43. No.1. P.120-128. (russian)
- [6] D.V. Belykh, E. V. Buravlev, P. A. Slepukhin, and A. V. Kuchin Atropoisomerism of 13-N,N-Dimethylamide-15,17-dimethyl Ester of Chlorin e6 from the Data of X-Ray, <sup>1</sup>H NMR, and HPLC. Russian Journal of General Chemistry. 2010. Vol.80. No.11. P.2382-2386. (russian)

## Full Paper

- [7] S.Y. Lee, N. Jagerovich, K.M. Smith. Use of the chlorophyll derivative purpurin 18 for synthesis for use in photodynamic therapy. J. Chem. Soc. Perkin Trans 1. 1993. P.2369-2377.
- [8] N.N. Lebedev. Chemistry and technology of basic organic and petrochemical synthesis. Ed. 2. Moscow: Chemistry. 1991. C.730. (russian)